



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# Comparison of upstream Te profiles with divertor heat flux and its implications on parallel and perpendicular transport in the SOL of DIII-D H-mode plasmas

M. A. Makowski, C. J. Lasnier, G. D. Porter, A. W. Leonard, J. A. Boedo, J. G. Watkins, D. N. Hill

June 17, 2010

37th European Physical Society Conference on Plasma  
Physics  
Dublin, Ireland  
June 21, 2010 through June 25, 2010

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

# Comparison of upstream $T_e$ profiles with divertor heat flux and its implications on parallel and perpendicular transport in the SOL of DIII-D H-mode plasmas

M.A. Makowski<sup>1</sup>, C.J. Lasnier<sup>1</sup>, G.D. Porter<sup>1</sup>, A.W. Leonard<sup>2</sup>, J.A. Boedo<sup>3</sup>,  
J.G. Watkins<sup>4</sup>, and D.N. Hill<sup>1</sup>

<sup>1</sup>*Lawrence Livermore National Laboratory, Livermore, California 94550, USA.*

<sup>2</sup>*General Atomics, P.O. Box 85608, San Diego, CA 92186-5608, USA.*

<sup>3</sup>*University of California-San Diego, La Jolla, California 92093, USA.*

<sup>4</sup>*Sandia National Laboratories, Albuquerque, New Mexico, USA*

## I. Introduction

On DIII-D we have performed a series of experiments designed to compare the upstream Thomson and midplane probe measurements of  $T_e$  with the downstream divertor heat flux width. We find that there is only a weak dependence of the heat flux width on the temperature gradient scale length, in contrast to the strong dependence predicted by simple two-point models [1]. The UEDGE code [2] has also been used to model a number of the discharges and we find that the flows significantly affect the heat flux profiles.

Correlations between the heat flux width and a wide variety of plasma parameters have been made with the result that the only dependence found is on the plasma current,  $I_p$ . No dependence was found on,  $P_{SOL}$ ,  $P_{inj}$ ,  $n_e$ ,  $B_T$ , the shear and normalized pressure gradient at the 95% flux surface ( $s_{95}$ ,  $\alpha_{95}$ ), nor collisionality. Comparing the DIII-D data with other multi-machine scaling relations, we find best agreement with the JET scaling [3].

## II. Comparison of Upstream and Midplane $T_e$ Profiles

Upstream Thomson profiles of  $T_e$  and  $n_e$  measured at the upper outer region of the plasma at a major radius of 1.94 m were mapped to the midplane for comparison with the heat flux profiles. The profiles and heat flux data were “edge localized mode (ELM) synchronized” using a method of conditionally averaging the data falling between ELMs over many ELM cycles. Independent exponential fits to the core- and scrape-off layer (SOL)-side data were made to obtain measurements of the profile gradient scale lengths as shown in Fig. 1. The core-side fits were clearly biased by the pedestal and not used in the subsequent analysis. SOL layer widths inferred from tanh-fits were also analyzed and showed the same trends as the SOL-side fits. To be consistent with other published data SOL-side gradient scale lengths were used in the following analysis.

A database was established by dividing the shot into 200 ms segments and averaging the various plasma parameters and profile data over this period. Additional parameters, such as

shear and normalized pressure gradient, were included in the database in order to perform regression analysis on them.

### III. Heat Flux Widths

Divertor temperature profiles were measured with an IRTV camera mounted on the top of the DIII-D vacuum vessel. The heat flux to the target plate was then inferred using the THEODOR [4] code. To arrive at a heat flux width, offset exponentials,  $a_0 + a_1 e^{x/\lambda}$ , were fit to both the left hand and right hand sides of the profiles. This was necessary as the baseline on the two sides of the profile were, in general, different. With this method an effective Loarte width (ratio of the integral of the heat flux profile to the peak heat flux) [5] could be calculated resulting in,

$$\lambda_q = \left( \lambda_{\text{left}} + \lambda_{\text{right}} \right) \frac{R_{\text{div}} B_{\theta}^{\text{div}}}{R_{\text{mp}} B_{\theta}^{\text{mp}}},$$

where the factor,  $R_{\text{div}} B_{\theta}^{\text{div}} / R_{\text{mp}} B_{\theta}^{\text{mp}}$ , accounts for the flux expansion between the target plate and midplane. Figure 2 shows a typical outer divertor heat flux profile mapped to the midplane with offset exponentials fit to both sides of the peak.

The heat flux widths were fit to power scalings of  $I_p$ ,  $P_{\text{SOL}}$ ,  $P_{\text{inj}}$ ,  $n_e$ ,  $B_T$ , shear and normalized pressure gradient at the 95% flux surface ( $s_{95}$ ,  $\alpha_{95}$ ) and collisionality. The only trend found was an inverse dependence on  $I_p$  as shown in Fig. 3. A weak dependence on  $B_T$  cannot be ruled out, as there were few good points in the data set at low fields ( $B_T < 1.5$  T).




Fig. 1. Upstream profiles of  $T_e$  and  $n_e$  mapped to the outer midplane. The SOL-side fit was used as a measure of the profile gradient scale length.




Fig. 2. Typical heat flux profile with offset exponentials fit to each side of the profile.




Fig. 3. Plot of the heat flux width,  $\lambda_q$ , versus  $I_p$  for two ranges of  $B_t$  (red and blue curve fits). Due to the weak dependence of  $\lambda_q$  on  $B_t$ , the two curves nearly overlap. The black line is fit to all the data (red and blue circles).

The measured heat flux width,  $\lambda_q$ , shows a very weak dependence on the upstream  $T_e$  gradient scale length,  $\lambda_{Te}$ , being essentially uncorrelated. This is shown in Fig. 4, which plots  $\lambda_q$  vs  $\lambda_{Te}$  and the fit to the data. This is in strong disagreement with the simple two-point model [1] that predicts  $\lambda_{Te} = (7/2) \lambda_q$ . This line is plotted in Fig. 4. Even with the poor correlation, this dependence can be ruled out. The observed weak dependence is not unreasonable as radial transport, SOL radiation, and divertor recycling affect the heat flux within flux tubes; effects not taken into account in the simple model.

#### IV. Scaling Relations

We have compared our results with several other scaling relations developed for the heat flux width. The first is the JET conduction limited scaling relation [3] given by

$$\lambda_q^{\text{JET}} (\text{mm}) = 2.41 \times 10^{-5} B_T^{-1} (\text{T}) P_{\text{SOL}}^{-1/2} (\text{MW}) n_e^{1/4} (\text{m}^{-3}) q_{95} R^2 (\text{m}) .$$

This is in quite good agreement with our data as shown in Fig. 5, which plots  $\lambda_q^{\text{DIII-D}}$  versus  $\lambda_q^{\text{JET}}$ . The bulk of the dependence derives from the variation in  $q_{95}/B_T \sim 1/I_p$ . Our data shows no dependence on  $n_e$ , while the JET scaling has a very weak  $n_e^{1/4}$  dependence, so there is little variation arising from this term. Our data shows no dependence on  $P_{\text{SOL}}$  though the JET scaling relation has a  $P_{\text{SOL}}^{1/2}$  dependence. Since  $R$  is a constant for our data, the remaining scaling relation essentially reduces to  $\lambda_q^{\text{JET}} \sim \lambda_q^{\text{JET}} \sim B_T^{-1} q_{95} \sim 1/I_p$ . This is quite close to the  $I_p^{-1.24}$  scaling found above (Fig. 3).

We have also considered two other multi-machine scaling relations [5]:

$$\lambda_q^{\text{H-1}} (\text{mm}) = 5.2 P_{\text{div}}^{0.44} (\text{MW}) B_T^{-0.45} (\text{T}) q_{95}^{0.57} ,$$

$$\lambda_q^{\text{H-2}} (\text{mm}) = 5.4 P^{0.38} (\text{MW}) B_T^{-0.71} (\text{T}) q_{95}^{0.30} .$$

These are in extremely poor agreement with the DIII-D data. The H-1 scaling predicts a dependence on plasma current of approximately  $I_p^{1/2}$  and no




Fig. 4. Plot of the heat flux width,  $\lambda_q$ , vs the Thomson electron profile gradient scale length in the scrape-off layer,  $\lambda_{Te, \text{SOL}}$ . Solid red line is a linear fit between the two parameters. The slope,  $s$ , is 1/10th that predicted by simple two-point models. Correlation coefficient = 0.124.




Fig. 5. Plot of  $\lambda_q^{\text{DIII-D}}$  vs  $\lambda_q^{\text{JET}}$  showing that the DIII-D data fits the JET scaling relation.

size dependence. The H-2 scaling predicts profile widths a factor of 10 lower than the measurements. In addition, there is no size dependence and there is no manifest dependence on  $I_p$ .

## V. UEDGE Simulations

Efforts are underway to model four representative points on the  $I_p$  scan of Fig. 3 with UEDGE [2] in order to determine what underlying physics might be changing with  $I_p$  to affect  $\lambda_q$ . Inputs to UEDGE are the power flux through the SOL and the midplane profiles of  $n_e$  and  $T_e$ . Transport coefficients are adjusted within UEDGE to obtain a match between the experimental and predicted profiles. Results are preliminary but still offer some insight.

Figure 6 shows a comparison of the measured heat flux profile and those obtained from a UEDGE simulation as a function of distance along the target plate for  $I_p = 1.5$  MA.

With impurities and drifts turned off and with impurities on and drifts off, the predicted profiles are narrow; approximately 50% of the width of the measured profile. Only with the flows (partially) turned on does the profile broaden by producing a shoulder on the SOL side. The profiles on the private flux side are very narrow and nearly constant for the varying conditions. The asymmetry between the private flux region and SOL is also reproduced by the simulation. The results appear to indicate that the plate physics and flows are important in determining the heat flux profile.

This work was supported by the U.S. Department of Energy under DE-AC52-07NA27344(LLNL), DE-FC02-04ER54698, DE-FG02-07ER54917, and DE-AC04-94AL85000.




Fig. 6. Heat flux profiles from UEDGE compared with measurement. Impurities have little effect on the profile whereas the flows have a pronounced effect.

- [1] C.S. Pitcher and P.C. Stangeby, Plasma Phys. Control. Fusion **39** (1997) 779.
- [2] T. Rognien, *et al.*, J. Nucl. Mater. **196-198** (1992) 347-351.
- [3] G. Kirnev, *et al.*, Plasma Phys. Control. Fusion **49** (2007) 689.
- [4] A. Hermann, *et al.*, Plasma Phys. Control. Fusion **37** (1995) 17.
- [5] A. Loarte, *et al.*, J. Nucl. Mater. **266-269** (1999) 587-592.